Children's Physician Group-Gynecology



Guidelines for referrals

Below is a list of guidelines to follow when referring a patient for a consultation to Children's Physician Group—Gynecology. These are meant to be general recommendations. If you have specific questions, call **404-785-DOCS (3627)** and ask to speak with the on-call gynecologist.

Common conditions treated

- Abnormal uterine bleeding
- Adnexal masses (ovarian/paraovarian cysts)
- Amenorrhea
- Complex contraception (pregnancy prevention in medically complex patients)
- Congenital adrenal hyperplasia
- Delayed puberty
- Disorders of sex development
- Dysmenorrhea
- Endometriosis
- Gender affirming care
- Hormone replacement therapy
- Menstrual suppression for special needs
- Mullerian (uterine) anomalies

- Pelvic inflammatory disease
- Precocious puberty
- Premature ovarian insufficiency
- Polycystic ovarian syndrome
- Urethral prolapse
- Vaginal anomalies
- Vulvar vaginal issues
 - Vaginal discharge
 - o Prepubertal vulvovaginitis
 - Labial/vulvar masses and ulcers
 - o Lichen sclerosus
 - Labial adhesions
 - Labial hypertrophy
 - o Genital tract trauma

For patients with the conditions listed below, we recommend a referral to our Adolescent Medicine Clinic, located at Hughes Spalding Hospital (phone: 404-785-9850), or an external gynecology provider.

- New patients >16 years: irregular periods, vulvovaginitis, contraceptive counseling, dysmenorrhea
- Return patients >16 years: controlled symptoms and no complex medical issues

Urgent referrals

Most issues we see do not warrant an urgent referral. However, if you feel your patient needs to be seen as soon as possible, note "urgent" on your referral. All referrals marked "urgent" are triaged to help make sure patients are seen in a timely fashion. If you wish to speak to the on-call pediatric gynecologist, call 404-785-9635. Generally, conditions that may warrant an urgent initial outpatient visit include, but are not limited to:

- Ovarian, pelvic, adnexal masses
- Acute genital tract trauma

- Differences of sexual development
- Vaginal/menstrual outflow tract obstruction

Routine referrals

The majority of conditions we see may not warrant an urgent evaluation given the available resources. These may include, but are not limited to, the following:

- Heavy menses
- Irregular or abnormal menstrual bleeding
- Painful menses (dysmenorrhea, endometriosis, pelvic pain NOS)
- Vaginal discharge or pain
- Precocious puberty
- Delayed puberty
- PCOS

Revised: March 21, 2022

Page 1 of 4

Referral checklist and guidelines for common diagnoses

When referring a patient for any reason, except transgender care, you must include office notes. Otherwise, we will <u>not</u> be able to schedule your patient. In the table below, we have listed the labs and/or documents we require for the most common referrals. If the suspected diagnosis is <u>not</u> listed below, you only need to include office notes and labs that have <u>already</u> been ordered.

Suspected diagnosis	Labs/documents required before scheduling	Criteria for referral	Suggested work-up	Possible initial management	
Heavy menses	☐ Office notes☐ Lab results if ordered	Bleeding > 7days, > 7 pads per day, menses resulting in anemia	 CBC Von Willebrand panel Fibrinogen TSH Iron studies 	***Aygestin; combined oral contraceptive pill (OCP); consider the risk for thrombosis before starting OCP*	
Precocious puberty	□ Office notes □ Growth curve □ Lab results if ordered	Breast, genital hair, vaginal bleeding prior to age 8	LHFSHEstradiolTSHProlactin		
Delayed puberty	□ Office notes□ Growth curve□ Lab results if ordered	No pubertal development by age 13	LHFSHEstradiolTSHProlactin		
Primary amenorrhea	☐ Office notes☐ Growth curve☐ Lab results ifordered	No menses by age 15 or 3 years after menarche	See irregular menses workup		
Irregular menses, oligomenorrhea, Polycystic Ovarian Syndrome (PCOS)	□ Office notes □ Growth curve □ Lab results if ordered	Irregular or absent bleeding Do labs if any androgenizing symptoms (acne, hirsutism)	 LH FSH Estradiol 17-hydroxy-progesterone Free testosterone DHEA-S TSH Fasting complete metabolic profile Fasting lipid profile hCG (urine or serum) Prolactin 	OCP is the first line of therapy; consider the risk for thrombosis before starting OCP* Metformin is used by some, but it is not an FDA-approved indication	
Pelvic mass	□ Office notes□ Imaging report*	We must have imaging report prior to scheduling appointment	 Patient to bring disc with images 		
Dysmenorrhea	□ Office notes □ Imaging if done			Ibuprofen 600mg TID; heating pads, warm bath, physical activity	
Complex contraception	☐ Office notes☐ Lab results if ordered	Patient has underlying medical problem that would prohibit adolescent medicine or general GYN from providing care	GonorrheaChlamydiaTrichomonas+/-RPR and HIV		

- *AUB labs should be drawn **before** starting hormone therapy, if indicated.
- ***Initial therapy in patient with heavy menstrual bleeding that is actively bleeding
 - Taper if hgb 8-11.9 and actively bleeding:
 - o Aygestin 10mg BID x3 days until 3 days after bleeding stops then continue 10mg daily OR
 - o Orthocyclen 1 tab q8 hours x3 days, then BID x2 days, then daily
 - Maintenance if hgb >11.9 or not actively bleeding:
 - o Aygestin 10mg daily OR
 - Orthocyclen 1 tab daily (may skip placebo week)
 - Send to Emergency Department for active bleeding (not spotting) and hgb <8

Growth curves

We require growth curves for <u>all</u> referred patients prior to scheduling. Note, it is very important to provide a **visual line graph**, ideally for both height and weight, although both are not required. Multiple points are preferred, *if available*. If you have only seen the patient once, we will accept graphs with single points.

Figure A






Figure B



Vitals with Age Percentiles	8/11/2016	12/15/2016	7/20/2017	4/20/2018	4/20/2018	6/20/2019	6/20/2019
Height percentile		84.1 %	79.1 %		60.5.%		86.9 %
Systolic percentile							
Diastolic percentile							
Weight percentile	38.2 %	39.4 %	32.7 %		61.3 %	1	42.6 %
Head Circumference percentile		98.4.%					
Length	*	95.3 cm	99.1 cm		102.cm		114.5 cm
Systolic			90		94		90
Diastolic			58		50		62
Head Circumference		20.250					
Pulse							
Weight	27 lb.	28 lb 6.1 oz	30-lb		36 lb 4 oz		39 lb 6.1 oz
Body Mass Index	,			15.8 kg/m2		13.63 kg/m2	
Body Mass Index percentile		5.7 %	4.3 %		65.0 %		6.8 %
BODY SURFACE AREA				0.68	1	0.75	



Office notes

Office notes are crucial in helping us determine the intricacies of your patient's case. Note, we require office notes *beyond* just the reason for referral. Whether you refer to them as Clinical Notes, History of Present Illness (HPI), Interval History or Notes, what we need are notes that the provider took during the last visit that explain the child's case and chief concern. A short "reason for referral" is **not** sufficient. See examples below.

Insufficient

Referral

Date requested: 08/16/2021

Requested by: FirstName LastName, NP **Referral to**: pediatric endocrinology

Summary of care provided:

Reason for referral/notes: breast buds and pubic hair

ICD code: Precocious puberty (ICD-10: E30.1)

Sufficient



PatientName is a XX-year-old female seen for follow-up visit via telemedicine for anxiety, depression and gender identity issues. Guardian called for crisis appointment as PatientName was distressed about breast development. Reports that she has never liked her body since age XX and identifies as a boy (symptoms worsened when she hit puberty). Patient would like to transition and talk about the process. Family is supportive.

Anxiety and depression: Overall mood has been stable. Not sleeping well, but she can focus on schoolwork. Denies self-harm or suicidal thoughts.

Insomia: Reports improved sleep with clonidine 0.1 mg at night.

Menarche: Cycles have been very heavy with clots, uses 5-6 pads/day. Cycles last 2-4 weeks.

Asthma: Mild persistent. On Ventolin HFA prn and Flovent daily. Needs refills.

