Cardiopulmonary resuscitation (CPR) for children with tracheostomy tubes



By knowing basic CPR, you can help your child during a heart or breathing emergency.

- If your child is usually responsive, make sure he is not responsive (tap him gently or shout).
- If unresponsive, send someone to call 911.

A – Open the AIRWAY

- Make sure the trach tube is not blocked with mucus, food or other objects.
- Suction the trach tube if needed.
- If the trach remains blocked, take it out and put in a new trach tube.

B₁ - Look, listen and feel for BREATHING

- Put your ear close to the trach tube.
- Watch his chest move look, listen and feel for breathing.

B₂ – Give 2 breaths with the resuscitator bag if your child is not getting enough air

• If you do not have the resuscitator bag, give 2 breaths mouth-to-trach tube to make the chest rise.

C - Begin CPR if your child shows no signs of life, such as coughing, moving or breathing

- Keep doing CPR until help arrives and they tell you to stop, until there are signs of life, or until 5 cycles have been completed.
- After 5 cycles of CPR, call 911 if they have not been called yet. Then, keep doing CPR. You can call sooner and while doing CPR if you have a cell phone with you at the time.
- If your child shows signs of life, stop CPR and check breathing.
- If not breathing, give rescue breaths until breathing begins or until help arrives.

Cardiopulmonary resuscitation for children with tracheostomy tubes, continued

	BABY Birth to 1 year	CHILD 1 year to puberty	ADULT Anyone past puberty
Chest compressions	100 to 120 per minute	100 to 120 per minute	100 to 120 per minute
Number of chest compressions and breaths	30 compressions and 2 breaths (each cycle) Breathe in gently , just enough to make the chest rise.	30 compressions and 2 breaths (each cycle) Breathe in just enough to make the chest rise.	30 compressions and 2 breaths (each cycle) Breathe in just enough to make the chest rise.
Chest compression depth	Push hard and fast (about 1/3 total depth of chest) – around 1 ½ inches. Allow chest to come all the way up each time.	Push hard and fast (about 1/3 total depth of chest) – around 2 inches. Allow chest to come all the way up each time.	Push hard and fast (about 1/3 total depth of chest) – at least 2 inches. Allow chest to come all the way up each time.
Hands	Use 2 fingers on breastbone just below nipple line. OR Use 2 thumbs (using encircling technique) on breastbone, just below nipple line. If unable to push down about 1/3 total	Use heel of 1 hand on breastbone between the nipples. For larger children, use 2 hands stacked on top of each other.	Use 2 hands stacked on top of each other - heel of the lower hand between the nipples. Point fingers in the same direction.
Rescue breathing (if heart is beating)	depth of chest, use heel of 1 hand instead. 1 breath every 3 to 5 seconds	You can always use 2 hands if 1 hand seems difficult. 1 breath every 3 to 5 seconds	1 breath every 5 to 6 seconds

This teaching sheet contains general information only. Talk with your child's doctor or a member of your child's healthcare team about specific care of your child.

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.