

Return to physical activity after a concussion

Lacrosse (boys)

Stage	Activity	Lacrosse exercises	Goal
1	<ul style="list-style-type: none"> Rest 	<ul style="list-style-type: none"> Rest other than everyday walking 	<ul style="list-style-type: none"> Promote recovery shortly after injury
2	<ul style="list-style-type: none"> Light aerobic activity 	<ul style="list-style-type: none"> 10 to 15 minutes of walking or stationary bike 	<ul style="list-style-type: none"> Add light aerobic activity Watch if symptoms return
3	<ul style="list-style-type: none"> Moderate aerobic activity Light resistance training 	<ul style="list-style-type: none"> 20 to 30 minutes of jogging with helmet and gloves Light weight lifting (1 set of 10 reps) 	<ul style="list-style-type: none"> Increase aerobic activity Watch if symptoms return
4	<ul style="list-style-type: none"> High intensity activity Non-contact lacrosse drills 	<ul style="list-style-type: none"> Sprints and agility drills Cradling, catching, scooping, fielding ground balls, shooting, change of direction, and give and go Waterfall drill, hamster drill, pinwheel drill and eagle eye drill Start with helmet and gloves. Move to full pads if symptom free 	<ul style="list-style-type: none"> Maximize aerobic activity Accelerate to full speed with change of directions (cuts) Start rotational head movements Watch if symptoms return
5	<ul style="list-style-type: none"> High intensity activity Limited contact lacrosse drills 	<ul style="list-style-type: none"> Sprints, agility drills and normal weight training Riding after the shot, riding off the end line, pick and roll, and 1 v 1 scramble No scrimmages Wear full pads 	<ul style="list-style-type: none"> Maximize aerobic activity Add high intensity strength training Add deceleration/rotational forces in controlled setting Watch if symptoms return
6	<ul style="list-style-type: none"> Full practice (after cleared by a provider) 	<ul style="list-style-type: none"> Normal training activities Normal scrimmages 	<ul style="list-style-type: none"> Frequent assessments during practice Watch if symptoms return
7	<ul style="list-style-type: none"> Return to play 	<ul style="list-style-type: none"> Normal game play 	<ul style="list-style-type: none"> Return to normal game play

Other guidelines:

If possible, a school athletic trainer or responsible adult should monitor the student as they move through each stage.

The student:

- May start Stage 2 when symptoms are improving and they can tolerate partial school days.
- May start Stage 3 when they can tolerate full school days with NO concussion symptoms.
- May move to next stage every 24 hours as long as symptoms do not return DURING or AFTER work out.
- May NOT move to Stage 6 without written clearance from a medical provider.
- Should return to their provider if they have trouble completing these stages without symptoms.

 choa.org/concussion

 404-785-KIDS (5437)

This is general information and is not specific medical advice. Always consult with a doctor or healthcare provider if you have any questions or concerns about the health of a child.

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.



Children'sSM
Healthcare of Atlanta