

Chamblee Primary Care ICD-10 documentation

Seven key impacts to ICD-10 documentation

1. Disease or disorder site
2. Acuity and/or encounter status of treatment
3. Etiology, causative agent, or disease type and injury/ poisoning cause, intent, activity at the time of the event and place event occurred
4. Underlying and associated conditions
5. Manifestations
6. Complications or adverse events
7. Supporting info such as lab values or socioeconomic key impacts to ICD-10 documentation

Most commonly used diagnoses

ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
Acute conjunctivitis, unspecified	372.00 372.00 372.00 372.00	H10.30	Unspecified acute conjunctivitis, unspecified eye	Refer to 7 key impacts to ICD-10 documentation
		H10.31	Unspecified acute conjunctivitis, right eye	
		H10.32	Unspecified acute conjunctivitis, left eye	
		H10.33	Unspecified acute conjunctivitis, bilateral	
Acute pharyngitis	462	J02.8	Acute pharyngitis due to other specified organisms	Upper Respiratory Disorders <ul style="list-style-type: none"> • Specify the acuity (i.e. acute, chronic, or recurrent) • Detail site (e.g. pharynx, tonsils, larynx, etc.) • Provide the infectious agent when known (e.g. streptococcus) • List any hypertrophy or obstruction • Document underlying or associated conditions (e.g. abscess) • Document any exposure to tobacco smoke
		J02.9	Acute pharyngitis, unspecified	
Acute upper respiratory infections of unspecified site	465.9	J06.9	Acute upper respiratory infection, unspecified	
Allergic rhinitis, cause unspecified	477.9	J30.0	Vasomotor rhinitis	
		J30.9	Allergic rhinitis, unspecified	
Streptococcal sore throat	034.0	J02.0	Streptococcal pharyngitis	
		J03.00	Acute streptococcal tonsillitis, unspecified	
		J03.01	Acute recurrent streptococcal tonsillitis	



ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
Acute suppurative otitis media without spontaneous rupture of ear drum	382.00	H66.001	Acute suppurative otitis media without spontaneous rupture of ear drum, right ear	<p>Otitis Media</p> <ul style="list-style-type: none"> Specify the acuity (i.e. acute, chronic, or recurrent) Detail the type (e.g. serous, mucoid, allergic, atticofurunculosis, tubotympanic, etc.) Differentiate non-suppurative and suppurative Identify laterality (right, left, bilateral) Report any spontaneous rupture of eardrum or myringitis State any related tobacco use, abuse, dependence, past history or smoke exposure (e.g. second hand) List any underlying disease (e.g. viral infection, influenza)
		H66.002	Acute suppurative otitis media without spontaneous rupture of ear drum, left ear	
		H66.003	Acute suppurative otitis media without spontaneous rupture of ear drum, bilateral	
		H66.004	Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, right ear	
		H66.005	Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, left ear	
		H66.006	Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, bilateral	
		H66.007	Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, unspecified ear	
		H66.009	Acute suppurative otitis media without spontaneous rupture of ear drum, unspecified ear	
Cough	786.2	R05	Cough	Refer to 7 key impacts to ICD-10 documentation
Fever, unspecified	780.60 780.60	R50.2	Drug induced fever	<p>Fever</p> <p>Identify the following:</p> <ul style="list-style-type: none"> When the origin is unknown Drug induced (identify drug) Underlying conditions and causes (e.g. neutropenic fever, heat, etc.) Post-procedure fever Post-vaccination fever Febrile nonhemolytic transfusion reaction
		R50.9	Fever, unspecified	
Health supervision for newborn under 8 days old	V20.31	Z00.110	Health examination for newborn under 8 days old	Refer to 7 key impacts to ICD-10 documentation
Morbid obesity	278.01	E66.01	Morbid (severe) obesity due to excess calories	

ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
------------	------------	-------------	-------------	--------------------

Need for prophylactic vaccination against Hemophilus influenza, type B [Hib]	V03.81	Z23	Encounter for immunization	
Need for prophylactic vaccination against Streptococcus pneumoniae [pneumococcus]	V03.82			
Need for prophylactic vaccination and inoculation against Diptheria-tetanus-pertussis, combined [DTP] [DTaP]	V06.1			
Need for prophylactic vaccination and inoculation against influenza	V04.81			
Need for prophylactic vaccination and inoculation against other combinations of diseases	V06.8			
Need for prophylactic vaccination and inoculation against other viral diseases	V04.89			
Need for prophylactic vaccination and inoculation against varicella	V05.4			
Need for prophylactic vaccination and inoculation against viral hepatitis	V05.3			
Need for prophylactic vaccination with diphtheria-tetanus-pertussis with poliomyelitis (DTP + polio) vaccine	V06.3			
Need for prophylactic vaccination with measles-mumps-rubella [MMR] vaccine	V06.4			

The ICD-10 Provider Specialty Tool Kit is meant to assist providers with documentation best practices necessary for the transition from ICD-9-CM to ICD-10-CM. This tool kit includes the most commonly used diagnoses for a specialty using 2012 claims data for top encounters and top number of dollars per diagnosis. The tool kit can be used as a quick reference guide for providers and can also be shared with Practice Managers. This tool kit is not meant to be a coding cross walk and therefore should not be used as such. Questions regarding this tool kit and/or ICD-10 should be submitted via email to ICD10@choa.org.