

Orthotics and Prosthetics ICD-10 documentation

Seven key impacts to ICD-10 documentation

1. Disease or disorder site
2. Acuity and/or encounter status of treatment
3. Etiology, causative agent, or disease type and injury/
poisoning cause, intent, activity at the time of the event
and place event occurred
4. Underlying and associated conditions
5. Manifestations
6. Complications or adverse events
7. Supporting info such as lab values or socioeconomic key
impacts to ICD-10 documentation

Most commonly used diagnoses

ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
Abnormality of gait	781.2	R26.0	Ataxic gait	Abnormality of gait Identify condition as: <ul style="list-style-type: none"> • Ataxic gait • Paralytic gait • Unsteadiness on feet • Other or unspecified abnormalities of gait and mobility
		R26.1	Paralytic gait	
		R26.81	Unsteadiness on feet	
		R26.89	Other abnormalities of gait and mobility	
		R26.9	Unspecified abnormalities of gait and mobility	
Congenital musculoskeletal deformities of skull, face, and jaw	754.0	Q67.0	Congenital facial asymmetry	Refer to 7 key elements of ICD-10 Documentation
Infantile cerebral palsy, unspecified	343.9	G80.9	Cerebral palsy, unspecified	
Intracranial injury of other and unspecified nature, without mention of open intracranial wound, with state of consciousness unspecified	854.00	S06.890A	Other specified intracranial injury without loss of consciousness, initial encounter	



ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
------------	------------	-------------	-------------	--------------------

		S06.890D	Other specified intracranial injury without loss of consciousness, subsequent encounter	
		S06.890S	Other specified intracranial injury without loss of consciousness, sequela	
Other congenital valgus deformities of feet	754.69	Q66.6	Other congenital valgus deformities of feet	
Scoliosis [and kyphoscoliosis], idiopathic	737.30	M41.112	Juvenile idiopathic scoliosis, cervical region	Scoliosis <ul style="list-style-type: none"> • Identify type of scoliosis: <ul style="list-style-type: none"> o Idiopathic o Infantile <ul style="list-style-type: none"> - Juvenile - Adolescent - Other o Secondary <ul style="list-style-type: none"> - Neuromuscular - Other o Thoracogenic o Other form o Unspecified • Identify site: <ul style="list-style-type: none"> o Cervical o Cervicothoracic o Thoracic o Thoracolumbar o Lumbar o Lumbosacral o Sacral/sacroccygeal (only applies to infantile idiopathic scoliosis) o Site unspecified
		M41.113	Juvenile idiopathic scoliosis, cervicothoracic region	
		M41.114	Juvenile idiopathic scoliosis, thoracic region	
		M41.115	Juvenile idiopathic scoliosis, thoracolumbar region	
		M41.116	Juvenile idiopathic scoliosis, lumbar region	
		M41.117	Juvenile idiopathic scoliosis, lumbosacral region	
		M41.119	Juvenile idiopathic scoliosis, site unspecified	
		M41.122	Adolescent idiopathic scoliosis, cervical region	
		M41.123	Adolescent idiopathic scoliosis, cervicothoracic region	
		M41.124	Adolescent idiopathic scoliosis, thoracic region	
		M41.125	Adolescent idiopathic scoliosis, thoracolumbar region	
		M41.126	Adolescent idiopathic scoliosis, lumbar region	
		M41.127	Adolescent idiopathic scoliosis, lumbosacral region	
		M41.129	Adolescent idiopathic scoliosis, site unspecified	
		M41.20	Other idiopathic scoliosis, site unspecified	
		M41.22	Other idiopathic scoliosis, cervical region	
		M41.23	Other idiopathic scoliosis, cervicothoracic region	
M41.24	Other idiopathic scoliosis, thoracic region			
M41.25	Other idiopathic scoliosis, thoracolumbar region			
M41.26	Other idiopathic scoliosis, lumbar region			

ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
		M41.27	Other idiopathic scoliosis, lumbosacral region	
Talipes, unspecified	754.70	Q66.89	Other specified congenital deformities of feet	Refer to 7 key elements of ICD-10 Documentation
Torticollis, unspecified	723.5	M43.6	Torticollis	
Traumatic amputation of leg(s) (complete) (partial), unilateral, below knee, without mention of complication	897.0	S88.111A	Complete traumatic amputation at level between knee and ankle, right lower leg, initial encounter	Traumatic amputation of leg <ul style="list-style-type: none"> • Identify type as complete or partial • Identify location and level of the amputation • Specialty laterality (right, left) • Identify type of encounter: initial, subsequent, sequela • Specify any complications
		S88.111D	Complete traumatic amputation at level between knee and ankle, right lower leg, subsequent encounter	
		S88.111S	Complete traumatic amputation at level between knee and ankle, right lower leg, sequela	
		S88.112A	Complete traumatic amputation at level between knee and ankle, left lower leg, initial encounter	
		S88.112D	Complete traumatic amputation at level between knee and ankle, left lower leg, subsequent encounter	
		S88.112S	Complete traumatic amputation at level between knee and ankle, left lower leg, sequela	
		S88.119A	Complete traumatic amputation at level between knee and ankle, unspecified lower leg, initial encounter	
		S88.119D	Complete traumatic amputation at level between knee and ankle, unspecified lower leg, subsequent encounter	
		S88.119S	Complete traumatic amputation at level between knee and ankle, unspecified lower leg, sequela	
		S88.121A	Partial traumatic amputation at level between knee and ankle, right lower leg, initial encounter	
		S88.121D	Partial traumatic amputation at level between knee and ankle, right lower leg, subsequent encounter	
		S88.121S	Partial traumatic amputation at level between knee and ankle, right lower leg, sequela	
		S88.122A	Partial traumatic amputation at level between knee and ankle, left lower leg, initial encounter	

ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
------------	------------	-------------	-------------	--------------------

		S88.122D	Partial traumatic amputation at level between knee and ankle, left lower leg, subsequent encounter	
		S88.122S	Partial traumatic amputation at level between knee and ankle, left lower leg, sequela	
		S88.129A	Partial traumatic amputation at level between knee and ankle, unspecified lower leg, initial encounter	
		S88.129D	Partial traumatic amputation at level between knee and ankle, unspecified lower leg, subsequent encounter	
		S88.129S	Partial traumatic amputation at level between knee and ankle, unspecified lower leg, sequela	

The ICD-10 Provider Specialty Tool Kit is meant to assist providers with documentation best practices necessary for the transition from ICD-9-CM to ICD-10-CM. This tool kit includes the most commonly used diagnoses for a specialty using 2012 claims data for top encounters and top number of dollars per diagnosis. The tool kit can be used as a quick reference guide for providers and can also be shared with Practice Managers. This tool kit is not meant to be a coding cross walk and therefore should not be used as such. Questions regarding this tool kit and/or ICD-10 should be submitted via email to ICD10@choa.org.