

Transplant Heart ICD-10 documentation

Seven key impacts to ICD-10 documentation

1. Disease or disorder site
2. Acuity and/or encounter status of treatment
3. Etiology, causative agent, or disease type and injury/
poisoning cause, intent, activity at the time of the event
and place event occurred
4. Underlying and associated conditions
5. Manifestations
6. Complications or adverse events
7. Supporting info such as lab values or socioeconomic key
impacts to ICD-10 documentation

Most commonly used diagnoses

ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
Aftercare following organ transplant	V58.44	Z48.21	Encounter for aftercare following heart transplant	Refer to 7 Key Impacts of ICD-10 Documentation
		Z48.22	Encounter for aftercare following kidney transplant	
		Z48.23	Encounter for aftercare following liver transplant	
		Z48.24	Encounter for aftercare following lung transplant	
		Z48.280	Encounter for aftercare following heart-lung transplant	
		Z48.288	Encounter for aftercare following multiple organ transplant	
		Z48.290	Encounter for aftercare following bone marrow transplant	
Hypoplastic left heart syndrome	746.7	Q23.4	Hypoplastic left heart syndrome	
Congestive heart failure, unspecified	428.0	I50.20	Unspecified systolic (congestive) heart failure	Heart Failure <ul style="list-style-type: none"> • Specify the acuity (i.e., acute, chronic, or acute on chronic) • Identify the type of failure (e.g., systolic, diastolic, combined) • List any relationship of hypertension and/or chronic kidney disease to heart failure • Identify the underlying cause (e.g., surgery)
		I50.21	Acute systolic (congestive) heart failure	
		I50.22	Chronic systolic (congestive) heart failure	
		I50.23	Acute on chronic systolic (congestive) heart failure	
		I50.30	Unspecified diastolic (congestive) heart failure	
I50.31	Acute diastolic (congestive) heart failure			



ICD-9 Name	ICD-9 Code	ICD-10 Code	ICD-10 Name	Documentation tips
		I50.32	Chronic diastolic (congestive) heart failure	
		I50.33	Acute on chronic diastolic (congestive) heart failure	
		I50.40	Unspecified combined systolic (congestive) and diastolic (congestive) heart failure	
		I50.41	Acute combined systolic (congestive) and diastolic (congestive) heart failure	
		I50.42	Chronic combined systolic (congestive) and diastolic (congestive) heart failure	
		I50.43	Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure	
		I50.9	Heart failure, unspecified	
Other primary cardiomyopathies	425.4	I42.0	Dilated cardiomyopathy	Cardiomyopathy <ul style="list-style-type: none"> Specify the type: <ul style="list-style-type: none"> Dilated/congestive Obstructive hypertrophic Non-obstructive hypertrophic Endomyocardial fibrosis/disease Due to a substance (e.g. alcohol) List cardiomyopathy seen in other diseases such as amyloidosis, gout, glycogen storage disease, and thyrotoxicosis
		I42.5	Other restrictive cardiomyopathy	
		I42.8	Other cardiomyopathies	
Unspecified essential hypertension	401.9	I10	Essential (primary) hypertension	Hypertension <ul style="list-style-type: none"> Identify type of hypertension as being: <ul style="list-style-type: none"> Essential Secondary Hypertensive heart disease Hypertensive chronic kidney disease <ul style="list-style-type: none"> List stage of chronic kidney disease Hypertensive heart & chronic kidney disease List any related smoke exposure (second hand)

The ICD-10 Provider Specialty Tool Kit is meant to assist providers with documentation best practices necessary for the transition from ICD-9-CM to ICD-10-CM. This tool kit includes the most commonly used diagnoses for a specialty using 2012 claims data for top encounters and top number of dollars per diagnosis. The tool kit can be used as a quick reference guide for providers and can also be shared with Practice Managers. This tool kit is not meant to be a coding cross walk and therefore should not be used as such. Questions regarding this tool kit and/or ICD-10 should be submitted via email to ICD10@choa.org.